

(19)  Canadian  
Intellectual Property  
Office

An Agency of  
Industry Canada

Office de la Propriété  
Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

(11) **CA 2 358 146**

(13) **A1**

(40) 19.03.2003

(43) 19.03.2003

(12)

(21) 2 358 146

(51) Int. Cl. 7:

**A61K 35/12, A61K 41/00,  
A61P 15/06, A61K 38/17**

(22) 19.09.2001

(71)

BOCKSCH, LEILA,  
201 - 119 King Street, LONDON, O1 (CA).  
FEHMI, TARIK,  
4353 Wellssborough  
Place, MISSISSAUGA, O1 (CA).  
KUSZNIERUK, KORNEL PATRICK,

48 - 50 Chapman Court, LONDON, O1 (CA).

(72)



BOCKSCH, LEILA (CA).  
FEHMI, TARIK (CA).  
KUSZNIERUK, KORNEL PATRICK (CA).

(54) METHODE D'INHIBITION DES AVORTEMENTS D'ORIGINE IMMUNOLOGIQUE

(54) A METHOD OF INHIBITING IMMUNOLOGICALLY MEDIATED ABORTIONS

(57)

The invention disclosed teaches methods of inhibiting recurrent spontaneous abortion in a mammal through administration of apoptotic leukocytes. Said leukocytes may be derived from maternal or paternal sources. Several methods of apoptotic induction may be used to attain anti-abortogenic effects.





Office de la Propriété  
Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office

An agency of  
Industry Canada

CA 2358146 A1 2003/03/19

(21) **2 358 146**

(12) **DEMANDE DE BREVET CANADIEN  
CANADIAN PATENT APPLICATION**

(13) **A1**

(22) Date de dépôt/Filing Date: 2001/09/19

(41) Mise à la disp. pub./Open to Public Insp.: 2003/03/19

(51) Cl.Int.<sup>7</sup>/Int.Cl.<sup>7</sup> A61K 35/12, A61K 38/17, A61P 15/06,  
A61K 41/00



(71) Demandeurs/Applicants:

BOCKSCH, LEILA, CA;  
FEHMI, TARIK, CA;  
KUSZNIERUK, KORNEL PATRICK, CA

(72) Inventeurs/Inventors:

BOCKSCH, LEILA, CA;  
FEHMI, TARIK, CA;  
KUSZNIERUK, KORNEL PATRICK, CA

(54) Titre : METHODE D'INHIBITION DES AVORTEMENTS D'ORIGINE IMMUNOLOGIQUE  
(54) Title: A METHOD OF INHIBITING IMMUNOLOGICALLY MEDIATED ABORTIONS



(57) Abrégé/Abstract:

The invention disclosed teaches methods of inhibiting recurrent spontaneous abortion in a mammal through administration of apoptotic leukocytes. Said leukocytes may be derived from maternal or paternal sources. Several methods of apoptotic induction may be used to attain anti-abortion effects.

Canada

<http://opic.gc.ca> • Ottawa-Hull K1A 0C9 • <http://cipo.gc.ca>

OPIC • CIPQ 191

OPIC



CIPQ

**A METHOD OF INHIBITING IMMUNOLOGICALLY MEDIATED  
ABORTIONS**

**ABSTRACT**

The invention disclosed teaches methods of inhibiting recurrent spontaneous abortion in a mammal through administration of apoptotic leukocytes. Said leukocytes may be derived from maternal or paternal sources. Several methods of apoptotic induction may be used to attain anti-abortogenic effects.

## DISCLOSURE

### Field of the Invention

The invention teaches methods of modulating the immune system of a mammal in order to prevent spontaneous abortion.

### Background

#### *Recurrent Spontaneous Abortion*

Recurrent Spontaneous Abortion (RSA) affects 1-2% of women in North America (1). RSA is believed to be caused by abnormal activation of Th1 cells, these cells are generally suppressed in pregnancy. Lin *et al* demonstrated successful murine pregnancy is associated with upregulated production of the Th2 cytokines IL-4, IL-5 and IL-10 in the fetoplacental unit (2). This was postulated to inhibit development of Th1 cells, cells associated with induction of inflammatory responses. The Th2 biasing of immune response by pregnancy can be seen in C57BL/6 mice infected with *Leishmania*. Normally C57BL/6 mice are resistant to such infection because of a Th1 bias. When these mice are pregnant, due to the Th2 biasing of pregnancy, infection with *Leishmania* is possible (3). Administration of Th1 cytokines to pregnant mice can increase the rate of abortion (4), whereas administration of Th2 cytokine suppresses it (5). Therapies which prevent RSA, such as paternal lymphocyte infusion are associated with augmentation of Th2 responses (6,7). Therefore it appears that successful pregnancy requires suppression of Th1 and promotion of Th2 immune responses.

Paternal lymphocyte infusion therapy was originally devised by Govallo based on the idea that RSA occurs due to improper recognition of paternal antigens by maternal immune system cells (8). Through immunization with paternal lymphocytes, the mother's immune system will be primed to recognize the

paternal antigens on the placenta, thus immune response will occur. However, Govallo believed that stimulation of antiplacental immune response by injection of paternal lymphocytes would not cause an inflammatory response, but an immune response which would actually help the placenta to grow. A parallel concept was introduced by Prehn in cancer immunology who showed that host antitumor immune response in some circumstance can promote tumor growth (9). Although some clinical trials using paternal lymphocyte infusion have reported a decreased number of RSA (10,11), a recent double blind study reported no difference between treated and control groups (12). Nevertheless this practice is still in use today.

Rational exists for the systemic administration of Th2 cytokines in women prone to RSA. Since cytokine like IL-10 can be safely administered, and prevent activation of Th1 immune responses (13). The problem with systemic cytokine therapy is cost associated, as well as the possibility of global immune suppression.

Thus there is a need for treatment of RSA in an economical, safe and effective manner.

#### *Apoptosis*

Apoptosis, in contrast to necrosis, is a type of cell death in which membrane integrity of the cell is kept, thus preventing leakage of intracellular organelles to the extracellular milieu (14). The process of apoptosis exists in order to rid the organism of unwanted cells without causing an inflammatory response. In contrast, cells dying by necrosis release various intracellular components causing inflammation and tissue damage. The concept of "danger signals" was originally proposed by Matzinger (15) with the idea that innate signals of danger exist, which instruct the body to launch immune response or to ignore the antigen. Under Matzinger's paradigm, danger is associated with tissue necrosis, whereas lack of danger occurs when cells die by apoptosis. This is exemplified by studies showing that intracellular danger signals are released by cells undergoing necrotic

but not apoptotic cell death. Such danger signals include free-floating heat shock proteins which can activate dendritic cells and macrophages (16, 17). Following with the idea that cells undergoing necrosis cause immune activation through the danger pathway, apoptotic death should inhibit activation of immune response. One method by which apoptotic cells inhibit immune response is through the production of immunosuppressive cytokines such as IL-10. In a study by Gao *et al*, T cells induced to undergo apoptosis secreted IL-10 protein, as well as increased production of IL-10 transcripts. In the same study it was demonstrated that apoptotic T cells can program macrophages to inhibit activation of pro-inflammatory Th1 immune responses (18). In another study, Voll *et al* demonstrated that apoptotic lymphocytes can suppress production of TNF- $\alpha$  and IL-12 from peripheral blood mononuclear cells activated with lipopolysaccharide (LPS) (19).

### *Conclusion*

In light of observations that paternal lymphocytes infusions seem to diminish spontaneous abortion, and that lymphocytes undergoing apoptosis inhibit inflammatory responses, we investigated whether transfer of paternal and maternal origin apoptotic lymphocytes into abortion-prone mice would protect the fetus from immunologically mediated destruction.

### Summary of the Invention

The disclosed invention teaches a method of inhibiting abortogenic immune responses through administration of treated blood cells into a pregnant mammal. More specifically, paternal or maternal leukocytes are treated with apoptotic stimuli after which they are introduced into a pregnant female. This process, we postulate, results in the inhibition of inflammatory immune responses, allowing for completion of pregnancy in situations under which the immune response would normally result in fetal destruction.

The invention discloses methods of preparing paternal or maternal leukocytes, in a manner which endows the cells with anti-inflammatory properties. Disclosed is the utility of these cells in inhibiting anti-fetal immune responses. The direct effect of administration of the treated leukocytes on the pregnant female, that is the anti-abortion effect, is also disclosed.

#### Brief Description of the Drawings

Figure I illustrates time-dependent apoptotic effects of serum starvation on murine leukocytes.

Figure II illustrates the dose-dependent antiabortogenic effects of infusing apoptotic paternal cells to a pregnant mother induced to undergo abortion by LPS administration.

Figure III illustrates the dose-dependent antiabortogenic effects of infusing apoptotic maternal cells to a pregnant mother induced to undergo abortion by LPS administration.

Figure IV illustrates the dose-dependent antiabortogenic effects of infusing paternal apoptotic cells into the CBA/J X DBA/2 model of spontaneous abortion.

Figure V illustrates the dose-dependent antiabortogenic effects of infusing maternal apoptotic cells into the CBA/J X DBA/2 model of spontaneous abortion.

#### Detailed Description of Preferred Embodiments

This invention teaches methods of preventing immunologically-mediated spontaneous abortions through administration of apoptotic lymphocytes. This effect, we believe is mediated through production of soluble and membrane bound components on apoptotic cells which suppress inflammatory immune responses.

It is these inflammatory responses which bring about the demise of the conceptus. In the data displayed apoptosis was achieved through serum starvation of the leukocytes. In other studies we performed similar antiabortogenic effects can be seen by injection of leukocytes in which apoptosis was induced through other methods such as irradiation or treatment with protein synthesis inhibitors. In agreement with previously published studies, the immunomodulatory effects of apoptotic leukocytes are independent of the method by which apoptosis was induced (18, 19).

One embodiment of the invention mends itself applicable to the current practice of immunizing pregnant women at risk of spontaneous abortion with paternal lymphocytes. Paternal leukocytes can be purified from whole blood of the father through the Ficoll™ technique or other methods known to one skilled in the art. Apoptosis can be induced in these cells through serum starvation or by other methods known such as irradiation, treatment with protein synthesis inhibitors, treatment with ozone gas, or treatment with hyperthermia. Quality of the apoptotic cells can be assessed by Annexin-V staining as well as propidium iodide in order to ensure a high percentage of apoptotic but not necrotic cells are administered. Injection of apoptotic cells into the mother can occur via direct intravenous administration using syringe and needle, or continuous drip administration. The same procedure can be carried out using leukocytes derived from the mother.

Detection of cytokines associated with spontaneous abortion such as interferon- $\gamma$  can serve as an immune marker to guide the number of apoptotic cells injected, as well as the frequency of injection.



## Examples

### Example I

#### *Induction of Apoptosis by Serum Starvation*

BALB/c mice were euthanized by carbon dioxide asphyxiation and spleens were collected under aseptic conditions. Spleens were disassociated with a wire mesh and erythrocytes were lysed with hypotonic buffer. Lysed erythrocytes were removed by centrifugation. Splenocytes were cultured at a concentration of  $10^5$  cells/well in serum-free RPMI media in 96 well plates in a total volume of 200  $\mu$ l. At day 0, 1, 2, 3, and 4 apoptosis was assessed using ApoAlert™ Annexin V-FITC apoptosis kit (Clontech, Palo Alto, CA) according to the manufacturer's instructions. As illustrated in Figure I a time dependent increase in apoptotic cells was observed.

### Example II

#### *Dose-Dependent Inhibition of LPS-Induced Spontaneous Abortion by Infusion of Apoptotic Paternal Leukocytes*

Pregnancy was induced in 8 week old BALB/c females by crossing with C57/B6 males. Day zero of pregnancy was established as the day copulatory plug was observed. On day 5 of pregnancy a single injection of 4  $\mu$ g lipopolysaccharide from *Escherichia coli* (LPS: Sigma Chemicals, St. Louis, MO), was administered in 100  $\mu$ l of saline through the intraperitoneal route. On day 5 an injection of apoptotic paternal (C57/B6) leukocytes was administered intravenously through the tail vein. Apoptotic leukocytes were induced by serum starvation for 4 days as described in Example I. Mice were sacrificed on day 12 of pregnancy. As shown in Figure II administration of fresh splenocytes (Untreated leukocytes) did not inhibit LPS-induced abortions whereas injection of apoptotic leukocytes induced a dose-dependent inhibition of abortions. Abortions were quantified as

percent resorption identified by morphological appearance. Each bar represents the average ( $\pm$ Standard Deviation) of 7 mice.

#### Example III

##### *Dose-Dependent Inhibition of LPS-Induced Spontaneous Abortion by Infusion of Apoptotic Maternal Leukocytes*

Pregnancy was induced in 8 week old BALB/c females by crossing with C57/B6 males. Day zero of pregnancy was established as the day copulatory plug was observed. On day 5 of pregnancy a single injection of 4  $\mu$ g lipopolysaccharide from *Escherichia coli* (LPS: Sigma Chemicals, St. Louis, MO), was administered in 100  $\mu$ l of saline through the intraperitoneal route. On day 5 an injection of apoptotic maternal (BALB/c) leukocytes was administered intravenously through the tail vein. Apoptotic leukocytes were induced by serum starvation for 4 days as described in Example I. Mice were sacrificed on day 12 of pregnancy. As shown in Figure III administration of fresh splenocytes (Untreated leukocytes) did not inhibit LPS-induced abortions whereas injection of apoptotic leukocytes induced a dose-dependent inhibition of abortions. Abortions were quantified as percent resorption identified by morphological appearance. Each bar represents the average ( $\pm$ Standard Deviation) of 7 mice.

#### Example IV

##### *Dose-Dependent Inhibition Spontaneous Abortion in The CBA/J X DBA/2 Model by Infusion of Apoptotic Paternal Lymphocytes.*

CBA/J female mice were bred with DBA/2. Day zero of pregnancy was established as the day copulatory plug was observed. On day 5 an injection of apoptotic paternal (DBA/2) leukocytes was administered intravenously through the tail vein. Apoptotic leukocytes were induced by serum starvation for 4 days

as described in Example I. Mice were sacrificed on day 12 of pregnancy. As shown in Figure IV administration of fresh splenocytes (Untreated leukocytes) did not inhibit LPS-induced abortions whereas injection of apoptotic leukocytes induced a dose-dependent inhibition of abortions. Abortions were quantified as percent resorption identified by morphological appearance. Each bar represents the average ( $\pm$ Standard Deviation) of 7 mice.

#### Example V

##### *Dose-Dependent Inhibition Spontaneous Abortion in The CBA/J X DBA/2 Model by Infusion of Apoptotic Maternal Lymphocytes.*

CBA/J female mice were bred with DBA/2. Day zero of pregnancy was established as the day copulatory plug was observed. On day 5 an injection of apoptotic maternal (CBA/J) leukocytes was administered intravenously through the tail vein. Apoptotic leukocytes were induced by serum starvation for 4 days as described in Example I. Mice were sacrificed on day 12 of pregnancy. As shown in Figure V administration of fresh splenocytes (Untreated leukocytes) did not inhibit LPS-induced abortions whereas injection of apoptotic leukocytes induced a dose-dependent inhibition of abortions. Abortions were quantified as percent resorption identified by morphological appearance. Each bar represents the average ( $\pm$ Standard Deviation) of 7 mice.

The invention discloses a method of inhibiting immunologically mediated abortions in a mammal. It is to be understood that the above examples are presented only for clarity and that the invention may take other embodiments as practiced by one skilled in the art.

## References:

1. Coulman CB, Page WP, McIntyre JA. Overview of recurrent pregnancy loss. *Immunological Obstetrics*. Norton & Co, New York London, pp 333-356, 1992.
2. Lin H, Mosmann TR, Guilbert L, Tuntipopipat S, Wegmann TG. Synthesis of T helper 2-type cytokines at the maternal-fetal interface. *J Immunol*. 1993 Nov 1;151(9):4562-73.
3. Krishnan L, Guilbert LJ, Russell AS, Wegmann TG, Mosmann TR, Belosevic M. Pregnancy impairs resistance of C57BL/6 mice to *Leishmania major* infection and causes decreased antigen-specific IFN-gamma response and increased production of T helper 2 cytokines. *J Immunol*. 1996 Jan 15;156(2):644-52.
4. Chaouat G, Menu E, Clark DA, Dy M, Minkowski M, Wegmann TG. Control of fetal survival in CBA x DBA/2 mice by lymphokine therapy. *J Reprod Fertil*. 1990 Jul;89(2):447-58.
5. Chaouat G, Assal Meliani A, Martal J, Raghupathy R, Elliot J, Mosmann T, Wegmann TG. IL-10 prevents naturally occurring fetal loss in the CBA x DBA/2 mating combination, and local defect in IL-10 production in this abortion-prone combination is corrected by in vivo injection of IFN-tau. *J Immunol*. 1995 May 1;154(9):4261-8.
6. Szpakowski A, Malinowski A, Glowacka E, Wilczynski JR, Kolasa D, Dynski M, Tchorzewski H, Zeman K, Szpakowski M. The influence of paternal lymphocyte immunization on the balance of Th1/Th2 type reactivity in women with unexplained recurrent spontaneous abortion. *Ginekol Pol*. 2000 Jun;71(6):586-92.

7. Hayakawa S, Karasaki-Suzuki M, Itoh T, Ishii M, Kanaeda T, Nagai N, Takahashi-Yamamoto N, Tochigi M, Chishima F, Fujii TK, Oyama J, Kitanaka S, Satoh K. Effects of paternal lymphocyte immunization on peripheral Th1/Th2 balance and TCR V beta and V gamma repertoire usage of patients with recurrent spontaneous abortions. *Am J Reprod Immunol.* 2000 Feb;43(2):107-15.
  
8. Govallo VI, Sidel'nikov VM. Immunization of pregnant women with allogenic lymphocytes from the husband as a method of preventing spontaneous abortion. *Akush Ginekol (Mosk).* 1983 Dec;(12):25-8.
  
9. Prehn RT. Immunostimulation of the lymphodependent phase of neoplastic growth. *J Natl Cancer Inst.* 1977 Oct;59(4):1043-9.
  
10. Gatenby PA, Cameron K, Simes RJ, Adelstein S, Bennett MJ, Jansen RP, Shearman RP, Stewart GJ, Whittle M, Doran TJ. Treatment of recurrent spontaneous abortion by immunization with paternal lymphocytes: results of a controlled trial. *Am J Reprod Immunol.* 1993 Mar;29(2):88-94.
  
11. Mowbray JF, Gibbings C, Liddell H, Reginald PW, Underwood JL, Beard RW. Controlled trial of treatment of recurrent spontaneous abortion by immunisation with paternal cells. *Lancet.* 1985 Apr 27;1(8435):941-3.
  
12. Ober C, Karrison T, Odem RR, Barnes RB, Branch DW, Stephenson MD, Baron B, Walker MA, Scott JR, Schreiber JR. Mononuclear-cell immunisation in prevention of recurrent miscarriages: a randomised trial. *Lancet.* 1999 Jul 31;354(9176):365-9.
  
13. Reich K, Garbe C, Blaschke V, Maurer C, Middel P, Westphal G, Lippert U, Neumann C. Response of psoriasis to interleukin-10 is associated with

suppression of cutaneous type 1 inflammation, downregulation of the epidermal interleukin-8/CXCR2 pathway and normalization of keratinocyte maturation. *J Invest Dermatol.* 2001 Feb;116(2):319-29.

14. Kerr, J. F. R., Wyllie, A. H. & Currie, A. R. Apoptosis: a basic biological phenomenon with widespread implications in tissue kinetics. *Br. J. Cancer* 1972, 26, 239-257.

15. Matzinger P. Tolerance, danger, and the extended family. *Annu Rev Immunol.* 1994;12:991-1045.

16. Basu S, Binder RJ, Suto R, Anderson KM, Srivastava PK. Necrotic but not apoptotic cell death releases heat shock proteins, which deliver a partial maturation signal to dendritic cells and activate the NF-kappa B pathway. *Int Immunol.* 2000 Nov;12(11):1539-46.

17. Basu S, Binder RJ, Ramalingam T, Srivastava PK. CD91 is a common receptor for heat shock proteins gp96, hsp90, hsp70, and calreticulin. *Immunity.* 2001 Mar;14(3):303-13.


18. Gao Y, Herndon JM, Zhang H, Griffith TS, Ferguson TA. Antiinflammatory effects of CD95 ligand (FasL)-induced apoptosis. *J Exp Med.* 1998 Sep 7;188(5):887-96.

19. Voll G, Zhong J, Block M Immunosuppressive aspects of apoptotic peripheral blood mononuclear cells. *Nature* 390:350, 1997

## **CLAIMS**


1. A method of preventing recurrent spontaneous abortion in a mammal through
  - (a) Withdrawing leukocytes from a donor
  - (b) Inducing apoptosis of the donor leukocytes
  - (c) Administration of the donor leukocytes to a female susceptible to RSA
2. The method of claim I in which leukocyte defines a purified population of neutrophils, monocytes, dendritic cells, T cells or B cells.
3. The method of claim I in which leukocyte defines a combination of neutrophils, monocytes, dendritic cells, T cells and B cells.
4. The method of claim I where apoptosis is induced through ultraviolet irradiation, X-irradiation,  $\gamma$ -irradiation, serum starvation or crosslinking of Fas.
5. The method of claim I where the apoptotic cells are used to prevent recurrent spontaneous abortion in a mammal.
6. The method of claim I where the donor can be maternal or paternal in reference to the pregnancy.


**Figure 1. Time-Dependent Induction of Murine  
Leukocyte Apoptosis Induced by Serum  
Starvation**





**Figure II. Dose-Dependent Inhibition of LPS-Induced Spontaneous Abortion by Infusion of Apoptotic Paternal Leukocytes**





**Figure IV. Dose-Dependent Inhibition  
Spontaneous Abortion in the CBA/J X DBA/2  
Model by Infusion of Apoptotic Paternal  
Lymphocytes.**

